



# Cover Crop Chart

**GROWTH CYCLE**

A = Annual  
B = Biennial  
P = Perennial

**RELATIVE WATER USE**

☾ = Low  
🔵 = Medium  
💧 = High

**PLANT ARCHITECTURE**

☿ = Upright  
✳ = Upright-Spreading  
⚡ = Prostrate

-----Cool Season-----

-----Warm Season-----

---Grass---		-----Broadleaf-----								---Grass---							
A	<b>Barley</b> ☾									A	<b>Pearl millet</b> ☾						
A	<b>Oat</b> ☾	A	<b>Phacelia</b> ☾									A	<b>Foxtail millet</b> ☾				
A/P	<b>Ryegrass</b> ☾	A	<b>Flax</b> ☾									A	<b>Buckwheat</b> ☾	A	<b>Proso millet</b> ☾		
-----Legumes-----																	
A	<b>Wheat</b> ☾	A	<b>Spinach</b> ✳	B	<b>Turnip</b> 💧 ✳	A	<b>Field pea</b> ☾	A	<b>Berseem clover</b> ☾	A/P	<b>Medic</b> ✳	A	<b>Chickpea</b> ✳	A	<b>Sunflower</b> ☾	A	<b>Sudan grass</b> ☾
A	<b>Cereal rye</b> ☾	A	<b>Kale</b> ✳	A	<b>Radish</b> 💧 ✳	A	<b>Lentil</b> ✳	B/P	<b>Red clover</b> ☾	P	<b>Birdsfoot trefoil</b> ⚡	A	<b>Cowpea</b> ✳	A	<b>Safflower</b> ☾	A	<b>Teff</b> ☾
A	<b>Triticale</b> ☾	A/B	<b>Canola</b> ✳	B	<b>Beet</b> 💧 ✳	A	<b>Lupin</b> ☾	P	<b>White clover</b> ☾	P	<b>Sainfoin</b> ☾	A	<b>Soybean</b> ✳	A	<b>Squash</b> ⚡	A	<b>Grain sorghum</b> ☾
A	<b>Annual fescue</b> ☾	A/P	<b>Mustard</b> ✳	A/B	<b>Carrot</b> 💧 ✳	A/B	<b>Vetch</b> ☾	A/B	<b>Sweetclover</b> ☾	P	<b>Alfalfa</b> ☾	A	<b>Mung bean</b> ✳	P	<b>Chicory</b> ✳	A	<b>Corn</b> ☾

# Cover Crop Chart

The Cover Crop Chart is produced and distributed by the staff of the Northern Great Plains Research Laboratory, Mandan, ND.

The Cover Crop Chart represents a compendium of information from multiple sources in the U.S. and Canada. Primary sources of information included the Midwest Cover Crops Council, USDA-SARE, USDA-NRCS PLANTS database, and relevant peer-reviewed journal articles. Designation of warm/cool season crops is based on prevalent growth habits rather than photosynthetic pathway. Ranges for seeding depth take into consideration moisture conditions at planting and variation in soil texture. Values for crude protein and C:N ratio assume homogenous samples of aboveground plant material unless stated otherwise. Information on specific crops is occasionally generalized, approximate, and/or incomplete and may not reflect performance in on-farm conditions. USDA-ARS makes no guarantee to the performance of specific crops based on information provided herein. Crop sequencing information for annual cropping systems may be found in the Crop Sequence Calculator, v. 3.1.

Content and data for crops were assembled by Mark Liebig and Holly Johnson with input from Dave Archer, Heather Dose, Marvin Hatzenbuehler, John Hendrickson, Robert Kolberg, Steve Merrill, Kristine Nichols, Delmer Schlenker, Marty Schmer, Eric Scholljegerdes, Don Tanaka, Cal Thorson, and Dawn Wetch. Chart design by Mark Liebig, Holly Johnson, and Jill Gunderson. Website updates by Lori Wanner. The Cover Crop Chart was generated with input from producers and technicians in the Area IV Soil Conservation Districts of North Dakota and NRCS staff at the Bismarck and Dickinson Field/Area Offices.

## • Useful cover crop resources:

- **Managing Cover Crops Profitably, 3<sup>rd</sup> Ed. Andy Clark (Ed.). Handbook Series Book 9, Sustainable Agriculture Network, Beltsville, MD.**
- **<http://www.mccc.msu.edu> = Midwest Cover Crops Council**
- **<http://www.sarep.ucdavis.edu> = SARE, University of California-Davis**
- **<http://www.plants.usda.gov> = USDA-NRCS, PLANTS Database**



Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotope, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD). The United States Department of Agriculture prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital and family status. To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence, S.W., Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer. Mention of trade or manufacturer names is provided for information only and does not constitute endorsement by USDA-ARS.



## For further information please contact:

Cover Crop Chart  
USDA-ARS Northern Great Plains Research Laboratory  
P.O.Box 459 Mandan, ND 58554-0459  
Voice: 701 667-3000 FAX: 701 667-3054.  
**[www.mandan.ars.usda.gov](http://www.mandan.ars.usda.gov)**




# Barley (*Hordeum vulgare* L.)

- Cool Season, grass
- Annual
- Upright plant architecture
- Low water use
- Good salinity tolerance
- Seeding depth:  $\frac{3}{4}$  – 2 inches
- Crude protein: hay 10-15%, grain 11-15%
- Benefits from arbuscular mycorrhizal associations
- Self pollinator (wind)
- Rated 'very good' at scavenging nitrogen from the soil



USDA-ARS, NRPRL

[Back to Cover Crop Chart](#)



Sustainable Agriculture Research & Education,  
University of California, Davis

# Oat (*Avena sativa* L.)

- Cool Season, grass
- Annual
- Upright plant architecture
- Medium water use
- Fair salinity tolerance
- Seeding depth: 1 – 2 inches
- Crude protein: hay 9-15%, grain 13-18%
- Will form arbuscular mycorrhizal associations
- Self pollinator (wind)
- Rated 'very good' at scavenging nitrogen from the soil

[Back to Cover Crop Chart](#)



Sustainable Agriculture Research & Education,  
University of California, Davis



Midwest Cover Crops Council

# Ryegrass (*Lolium sp.*)

- Cool Season, grass
- Annual or perennial
- Upright plant architecture
- Major types:
  - Annual (Oregon, Italian, Australian, Common)
  - Perennial (English)
- Medium water use
- Fair salinity tolerance
- Seeding depth:  $\frac{1}{4}$  –  $\frac{1}{2}$  inch
- C:N ratio: 14 – 40
- Will form arbuscular mycorrhizal associations
- Self pollinator (wind)
- Rated 'very good' at scavenging nitrogen from the soil



Midwest Cover Crops Council

[Back to Cover Crop Chart](#)



# Wheat (*Triticum aestivum* L.)

- Cool season, grass
- Annual
- Upright plant architecture
- Includes spring and winter wheat varieties
- Medium water use
- Good to fair salinity tolerance
- Seeding depth: ½ – 1 ½ inches
- Crude protein: straw 4-10%, grain 12-16%
- C:N ratio: leaf 15-29, stem 31-65, root 24-74 , straw 80-95 [*end of season*]
- Benefits from arbuscular mycorrhizal associations
- Self pollinator (wind)
- Rated 'very good' at scavenging nitrogen from the soil

[Back to Cover Crop Chart](#)



USDA, NRCS, PLANTS Database



USDA-ARS, NPGRL

# Cereal rye (*Secale cereale* L.)

- Cool Season, grass
- Annual
- Upright plant architecture
- High water use
- Good salinity tolerance
- Seeding depth:  $\frac{1}{4}$  – 2 inches
- Crude protein: straw 4%, grain 14%
- C:N ratio: 40 – 48
- Will form arbuscular mycorrhizal associations
- Self pollinator (wind)
- Rated 'very good' at scavenging nitrogen from the soil



Midwest Cover Crops Council

[Back to Cover Crop Chart](#)

# Triticale

(*Triticale hexaploide* Lart.; *Triticosecale rimpaii* Wittm.)

- Cool Season, grass
- Annual
- Upright plant architecture
- Fall and spring types available
- High water use
- Good salinity tolerance
- Seeding depth: 1 ½ – 2 inches
- Crude protein: hay 9-16%, grain 17%
- Will form arbuscular mycorrhizal associations
- Self pollinator (wind)



[Back to Cover Crop Chart](#)

Midwest Cover Crops Council



# Annual fescue

(*Fetuca* sp.; *Vulpia myuros* L.)

- Cool Season, grass
- Annual
- Upright plant architecture
- Common names: Rattail fescue, Foxtail fescue
- Seeding depth: ¼ – 1 inch
- Crude protein: hay 8-10%
- Benefits from arbuscular mycorrhizal associations

[Back to Cover Crop Chart](#)



# Phacelia (*Phacelia tanacetifolia* Benth.)

- Cool Season, broadleaf
- Annual
- Upright plant architecture
- Low water use
- Low salinity tolerance
- Seeding depth:  $\frac{1}{8}$  –  $\frac{1}{4}$  inch
- C:N ratio: 10 – 15
- Grown as a nitrogen catch crop
- Attracts beneficial insects
- Will form arbuscular mycorrhizal associations



USDA, NRCS, PLANTS Database

[Back to Cover Crop Chart](#)

# Flax (*Linum usitatissimum*)

- Cool Season, broadleaf
- Annual
- Upright plant architecture
- Medium water use
- Fair salinity tolerance
- Seeding depth: ½ – 1 ½ inches
- Benefits from arbuscular mycorrhizal associations
- Flowers attract pollinators

[Back to Cover Crop Chart](#)



USDA-ARS, NRPRL

# Spinach (*Spinacia oleracea* L.)

- Cool Season, broadleaf
- Annual
- Upright and spreading plant architecture
- Low to medium water use
- Poor salinity tolerance
- Seeding depth:  $\frac{1}{4}$  –  $\frac{1}{2}$  inch
- Crude protein:  $\approx 20\%$
- C:N ratio: 6 – 8
- Sensitive to acid soils
- Belongs to the family Chenopodiaceae, which does not form arbuscular mycorrhizal associations



USDA, NRCS, PLANTS Database

[Back to Cover Crop Chart](#)



# Kale (*Brassica napus* L. var. *pabularia*)

- Cool Season, broadleaf
- Annual
- Upright and spreading plant architecture
- Major types: Siberian, Russian
- Kales belong to same genus and species as canola
- Medium water use
- Fair salinity tolerance
- Seeding depth:  $\frac{1}{4}$  –  $\frac{1}{2}$  inch
- Crude protein:  $\approx 30\%$
- C:N ratio: 10 – 30
- Does not form arbuscular mycorrhizal associations



Seed Ambassadors

[Back to Cover Crop Chart](#)



# Canola (*Brassica napus*)

- Cool Season, broadleaf
- Major types:
  - Annual (spring-type)
  - Biennial (winter-type)
- Upright and spreading plant architecture
- Also referred to as rapeseed
- Medium water use
- Good salinity tolerance
- Seeding depth:  $\frac{1}{4}$  – 1 inch
- Crude protein: hay 16%, grain 21%, silage 12%, pasture 17%
- C:N ratio: leaf 12-16, stem 21-37, root 24-43
- Does not form arbuscular mycorrhizal associations
- Rated 'very good' at scavenging nitrogen from the soil
- Flowers attract pollinators



USDA-ARS, NPGRL



Canola Council of Canada

# Mustard (*Brassica sp. L.*)

- Cool Season, broadleaf
- Annual or perennial
- Upright and spreading plant architecture
- Major types: brown, yellow, Indian, Oriental, etc.
- Low water use
- Poor salinity tolerance
- Seeding depth:  $\frac{1}{4}$  –  $\frac{1}{2}$  inch
- Crude protein: hay 10%, grain 24-35%
- C:N ratio: 10 – 30
- Mustard is related to crambe which has a low water use requirement
- Does not form arbuscular mycorrhizal associations
- Rated 'good' at scavenging nitrogen from the soil
- Plants from the *Brassica* group have potential to release compounds or metabolic by-products that work as bio-toxins against bacteria, fungi, insects, nematodes, and weeds
- Flowers may attract honeybees, lygus bugs, and hoverflies



Midwest Cover Crops Council



Crambe: USDA-ARS,  
NGPRL

[Back to Cover Crop Chart](#)

# Turnip (*Brassica rapa* L. var. *rapa*)

- Cool Season, broadleaf
- Biennial
- Upright and spreading plant architecture
- Root crop
- High water use
- Poor salinity tolerance
- Seeding depth:  $\frac{1}{4}$  –  $\frac{1}{2}$  inch
- Crude protein: tops 16%, root 12-14%
- Closely related to rutabagas
- Does not form arbuscular mycorrhizal associations
- Rated 'good' at scavenging nitrogen from the soil
- Flowers attract pollinators



Bonnie Plants



AMPAC Seed Company



Mas du Diable, January 2009

[Back to Cover Crop Chart](#)



# Radish (*Raphanus sativus*)

- Cool Season, broadleaf
- Annual
- Upright and spreading plant architecture
- Root crop
- Major types:
  - Oilseed (var. *oleiformis*)
  - Forage (var. *niger*)
- High water use
- Poor salinity tolerance
- Seeding depth:  $\frac{1}{4}$  –  $\frac{1}{2}$  inch
- Crude protein: 26-30%
- C:N ratio: oilseed 19 – 20
- Does not form arbuscular mycorrhizal associations
- Rated 'very good' at scavenging nitrogen from the soil
- Flowers attract pollinators



University of Maryland Extension, Fact Sheet 824

# Beet (*Beta vulgaris*)

- Cool Season, broadleaf
- Biennial
- Upright and spreading plant architecture
- Root crop
- High water use
- Variable salinity tolerance, depending on beet type.
- Seeding depth:  $\frac{1}{2}$  -  $\frac{3}{4}$  inch
- Crude protein: tops 12-15%, root 7-10%
- C:N ratio: tops 11 – 14
- Does not form arbuscular mycorrhizal associations
- Rated 'good' at scavenging nitrogen from the soil
- Self pollinator (wind)



Augustus Oils Limited



katynally.wordpress.com

[Back to Cover Crop Chart](#)



# Carrot (*Daucus carota* var. *sativus* L.)

- Cool Season, broadleaf
- Major types:
  - Biennial (cultivated)
  - Annual (wild)
- Upright and spreading plant architecture
- Root crop
- High water use
- Seeding depth:  $\frac{1}{8}$  -  $\frac{1}{4}$  inch
- Crude protein: 10%
- Forms arbuscular mycorrhizal associations
- Plants may bolt and flower starting in second year of growth. Flowers may attract honeybees.



[Back to Cover Crop Chart](#)

# Field pea (*Pisum sativum arvense* L.)

- Cool Season, broadleaf
- Annual
- Legume (N fixation)
- Upright plant architecture (vine)
- Low water use
- Poor salinity tolerance
- Seeding depth: 1 – 3 inches
- Crude protein: hay 14%, grain 24%, silage 15%
- C:N ratio: leaf 13-25, stem 27-83, root 17-27
- Forms arbuscular mycorrhizal associations
- Flowers attract bees

[Back to Cover Crop Chart](#)



Midwest Cover Crops Council



USDA-ARS, NGRPL

# Lentil (*Lens culinaris* Medik.)

- Cool Season, broadleaf
- Annual
- Legume (N-fixation)
- Upright and spreading plant architecture
- Low water use
- Poor salinity tolerance
- Seeding depth: 1 – 1 ½ inches
- Crude protein: hay 14%, grain 28%, silage 15%
- C:N ratio: leaf 11-21, stem 25-49, root 22-30
- Forms arbuscular mycorrhizal associations
- Self pollinated but flowers may attract pollinators



USDA-ARS, NPGRL

[Back to Cover Crop Chart](#)



# Lupin (*Lupinus sp.* L.)

- Cool Season, broadleaf
- Annual
- Legume (N-fixation)
- Upright plant architecture
- Examples include blue, narrow-leaved, European yellow, white, Spanish, etc.
- Low water use
- Prefers acid soils
- Seeding depth: 1 – 2 inches
- Crude protein: silage 15%
- C:N ratio: leaf 12-30, stem 25-49
- Does not form arbuscular mycorrhizal associations
- Flowers attract bees



USDA, NRCS, PLANTS Database



Sustainable Agriculture Research & Education,  
University of California, Davis

[Back to Cover Crop Chart](#)

# Vetch (*Vicia sp.*)

- Cool Season, broadleaf
- Annual or biennial
- Legume (N-fixation)
- Prostrate plant architecture (vine)
- Examples include common, hairy, purple, smooth, etc.
- Low to medium water use
- Poor salinity tolerance
- Seeding depth: 1 ½ – 2 ½ inches
- Crude protein: 13-20%
- C:N ratio: 10 – 19
- Forms arbuscular mycorrhizal associations
- Attracts pollinators



[Back to Cover Crop Chart](#)



# Berseem clover

## (*Trifolium alexandrinum* L.)

- Cool Season, broadleaf
- Annual
- Legume (N-fixation)
- Upright plant architecture
- Common name: Egyptian clover
- Low water use
- Fair salinity tolerance
- Seeding depth: ¼ – 1 inch
- Crude protein: 27-29%
- C:N ratio: 18 – 23
- Forms arbuscular mycorrhizal associations
- Flowers attract bees

[Back to Cover Crop Chart](#)



Wild Flowers of Israel, *Trifolium alexandrinum*



Sustainable Agriculture Research & Education,  
University of California, Davis

# Red clover (*Trifolium pratense* L.)

- Cool Season, broadleaf
- Biennial; short-lived perennial
- Legume (N-fixation)
- Upright plant architecture
- Common names: medium red clover, mammoth clover
- Medium water use
- Poor salinity tolerance
- Seeding depth: ¼ – ½ inch
- Crude protein: hay 15%
- C:N ratio: 15 – 23
- Forms arbuscular mycorrhizal associations
- Flowers attract bees



[Back to Cover Crop Chart](#)

# White clover (*Trifolium repens* L.)

- Cool Season, broadleaf
- Perennial
- Legume (N-fixation)
- Upright plant architecture
- Common names: Ladino, Dutch white, New Zealand white
- Medium water use
- Poor salinity tolerance
- Seeding depth: ¼ inch
- Crude protein: 24-30%
- C:N ratio: 13 – 23
- Forms arbuscular mycorrhizal associations
- Flowers attract bees



Sustainable Agriculture Research & Education,  
University of California, Davis



[Back to Cover Crop Chart](#)



# Sweetclover (*Melilotus* sp. L)

- Cool Season, broadleaf
- Annual or biennial
- Legume (N-fixation)
- Two varieties
  - yellow *Melilotus officinalis* L.
  - white *Melilotus alba* L.
- Upright plant architecture
- Moderate water use
- Fair salinity tolerance
- Seeding depth: ½ inch
- Crude protein: hay 11-18%
- C:N ratio: 12 – 23
- Forms arbuscular mycorrhizal associations
- Attracts pollinators



WHITE VARIETY  
Midwest Cover Crops Council



YELLOW VARIETY  
Midwest Cover Crops Council





# Medic (*Medicago* spp.)

- Cool Season, broadleaf
- Annual or perennial
- Legume (N-fixation)
- Upright and spreading plant architecture
- Over 35 known species of medics exist. Examples include barrel, black, burr, etc.
- Low water use
- Poor to fair salinity tolerance
- Seeding depth: ¼ inch
- Crude protein: black medic 19-21%
- Forms arbuscular mycorrhizal associations
- Attracts pollinators



BLACK MEDIC  
Midwest Cover Crops Council



BURR MEDIC  
Sustainable Agriculture Research & Education,  
University of California, Davis

[Back to Cover Crop Chart](#)

# Birdsfoot trefoil (*Lotus corniculatus*)

- Cool Season, broadleaf, legume
- Perennial
- Legume (N-fixation)
- Prostrate plant architecture
- Low to medium water use
- Fair salinity tolerance
- Seeding depth:  $\frac{1}{4}$  –  $\frac{1}{2}$  inch
- Crude protein: hay 16%
- Forms arbuscular mycorrhizal associations
- Attracts pollinators



USDA, NRCS, PLANTS Database

[Back to Cover Crop Chart](#)



Sustainable Agriculture Research & Education,  
University of California, Davis

# Sainfoin (*Onobrychis viciifolia* Scop.)

- Cool Season, broadleaf
- Perennial
- Legume (N-fixation)
- Upright plant architecture
- Medium to high water use
- Fair salinity tolerance
- Seeding depth:  $\frac{1}{4}$  –  $\frac{3}{4}$  inch
- Crude protein: hay 13-20%
- Forms arbuscular mycorrhizal associations
- Attracts pollinators



Forestry Images

[Back to Cover Crop Chart](#)

# Alfalfa (*Medicago sativa* L.)

- Cool Season, broadleaf
- Perennial
- Legume (N-fixation)
- Upright plant architecture
- High water use
- Poor salinity tolerance
- Seeding depth:  $\frac{1}{4}$  –  $\frac{1}{2}$  inch
- Crude protein: hay or silage 14-22%
- C:N ratio: 11 – 13
- Non-dormant cultivars can perform like an annual
- Forms arbuscular mycorrhizal associations
- Good at scavenging nitrogen from the soil
- Attracts pollinators



Midwest Cover Crops Council

[Back to Cover Crop Chart](#)



# Chickpea (*Cicer arietinum* L.)

- Warm Season, broadleaf
- Annual
- Legume (N-fixation)
- Two types
  - Desi
  - Kabuli
- Upright and spreading plant architecture
- Common name: garbanzo bean
- Low water use
- Poor salinity tolerance
- Seeding depth: 1 ½ – 2 inches
- Crude protein: straw 6%, grain 22%
- C:N ratio: leaf 10-15, stem 26-56, root 16-27
- Forms arbuscular mycorrhizal associations
- Flowers attract bees



[www.mathildasanthropologyblog.wordpress.com](http://www.mathildasanthropologyblog.wordpress.com)



Forestry Images


# Cowpea (*Vigna unguiculata* L.)

- Warm Season, broadleaf
- Annual
- Legume (N-fixation)
- Upright and spreading plant architecture (vine)
- Alternate names: Southern pea, Black-eye pea
- Low water use
- Poor salinity tolerance
- Seeding depth:  $\frac{3}{4}$  – 1 inch
- Crude protein: 19-24%
- C:N ratio: 18 – 22
- Forms arbuscular mycorrhizal associations
- Attracts pollinators

[Back to Cover Crop Chart](#)



USDA-ARS, NGPRL, dry bean



# Soybean (*Glycine max* (L.) Merr.)

- Warm Season, broadleaf
- Annual
- Legume (N-fixation)
- Upright and spreading plant architecture
- Medium water use
- Poor salinity tolerance
- Seeding depth: 1 – 2 inches
- Crude protein: hay 17%, grain 42%
- C:N ratio: leaf 14, stem 39, root 34
- Forms arbuscular mycorrhizal associations
- Self pollinated but flowers may attract pollinators

[Back to Cover Crop Chart](#)



USDA-ARS, NGRPL



Midwest Cover Crops Council



# Mung bean (*Vigna radiata* L.)

- Warm Season, broadleaf
- Annual
- Legume (N-fixation)
- Upright and spreading plant architecture
- Low to medium water use
- Poor salinity tolerance
- Seeding depth: 1 ½ – 3 inches
- Crude protein: 16-23%
- C:N ratio: 10 – 15
- Forms arbuscular mycorrhizal associations
- Self pollinated

[Back to Cover Crop Chart](#)



Top Crop Manager



# Amaranth (*Amaranthus sp.*)

- Warm Season, broadleaf
- Annual
- Upright plant architecture
- Over 50 species
- Low water use
- Tolerant of heat and drought
- Seeding depth: ½ – 2 inches
- Crude protein: ≈14%
- In-vitro dry matter digestibility: ≈ 71%
- Does not form arbuscular mycorrhizal associations
- Self pollinated (wind) but flowers may attract pollinators

[Back to Cover Crop Chart](#)



USDA-ARS, Germplasm Resources Information Network  
\*additional images at [ForestryImages.com](http://ForestryImages.com)

# Buckwheat

(*Fagopyrum esculentum* Moench; *Fagopyrum sagittatum* Gilib)

- Cool Season, broadleaf
- Warm season growth characteristics
- Annual
- Upright plant architecture
- Medium water use
- Poor salinity tolerance
- Enhances soil P availability
- Seeding depth: ½ inch
- Crude protein: straw 5%, grain 13%
- C:N ratio: leaf 8-10, stem 12-32, root 28-47
- Does not form arbuscular mycorrhizal associations
- Attracts pollinators



USDA-ARS, NPGRL

[Back to Cover Crop Chart](#)

# Sunflower (*Helianthus annuus* L.)

- Warm season growth characteristics, broadleaf
- Annual
- Upright plant architecture
- High water use
- Fair salinity tolerance
- Deep rooted
- Effective at 'mining' mobile nutrients deep in the soil profile
- Seeding depth: 1 – 3 ½ inches
- Crude protein: silage 11-12%, grain 20-28%
- C:N ratio: leaf 11-14, stem 41-46, root 50-68, flower 14-19 [*end of season*]
- Will form arbuscular mycorrhizal associations
- Flowers attract pollinators



USDA-ARS, NPGRL



GAP Photos



# Safflower (*Carthamus tinctorius* L.)

- Warm Season, broadleaf
- Annual
- Upright plant architecture
- High water use
- Good salinity tolerance
- Deep rooted
- Effective at 'mining' mobile nutrients deep in the soil profile
- Seeding depth: 1 – 1 ½ inches
- Crude protein: hay 10-13%, grain 18%
- C:N ratio: leaf 21, stem 56, root 73
- Will form arbuscular mycorrhizal associations
- Flowers attract pollinators



Forestry Images



USDA-ARS, NPGRL

[Back to Cover Crop Chart](#)



# Squash (*Cucurbita* sp.)

- Warm Season, broadleaf
- Annual
- Prostrate plant architecture (vine)
- Examples include butternut, winter, crookneck, and zucchini (summer squash)
- Genus encompasses summer/winter squash, pumpkin, and some gourds
- Seeding depth: ½ – 1 inch
- Forms arbuscular mycorrhizal associations
- Attracts pollinators

[Back to Cover Crop Chart](#)



# Chicory (*Cichorium intybus* L.)

- Warm Season, broadleaf
- Perennial
- Upright and spreading plant architecture (vine)
- Seeding depth:  $\frac{1}{8}$  –  $\frac{1}{2}$  inch
- Protein levels: 10-32% depending on plant maturity
- Will form arbuscular mycorrhizal associations
- Attracts pollinators

[Back to Cover Crop Chart](#)



Forestry Images

# Pearl millet (*Pennisetum glaucum* L.)

- Warm Season, grass
- Annual
- Upright plant architecture
- Low water use
- Poor salinity tolerance
- Seeding depth: ½ – 1 inch
- Crude protein: hay 13%
- Will form arbuscular mycorrhizal associations
- Self pollinator (wind)



Midwest Cover Crops Council



[Back to Cover Crop Chart](#)



Forestry Images



# Foxtail millet (*Setaria italica* L.)

- Warm Season, grass
- Annual
- Upright plant architecture
- Low water use
- Poor salinity tolerance
- Seeding depth: 1 inch
- Crude protein: hay 15%
- Will form arbuscular mycorrhizal associations
- Self pollinator (wind)



[Back to Cover Crop Chart](#)



# Proso millet (*Panicum milaceum* L.)

- Warm Season, grass
- Annual
- Upright plant architecture
- Medium water use
- Poor salinity tolerance
- Seeding depth: 1 inch
- Crude protein: hay 10%
- C:N ratio: leaf 12-16, stem 12-35, root 17-26
- Will form arbuscular mycorrhizal associations
- Self pollinator (wind)



USDA-ARS, NRPRL

[Back to Cover Crop Chart](#)

# Sudan grass

## *(Sorghum bicolor* L. Moench)

- Warm Season, grass
- Annual
- Upright plant architecture
- Common names: Sudan grass, Sorghum
  - *Note: These were formerly separate species that have been combined*
- Medium water use
- Fair salinity tolerance
- Seeding depth: 1 inch
- Crude protein: hay 7-11%, silage 6-17%
- C:N ratio: 48 - 63
- Benefits from arbuscular mycorrhizal associations
- Self pollinator (wind)
- Stress conditions that limit growth (e.g., drought, frost) can contribute to prussic acid accumulation in leaves



Forestry Images



[Back to Cover Crop Chart](#)

# Teff

## *(Eragrostis tef (Zuccagni) Trotter)*

- Warm Season, grass
- Annual
- Upright plant architecture
- Medium water use
- Poor salinity tolerance
- Seeding depth: ½ inch
- Crude protein: 10-18%
- Benefits from arbuscular mycorrhizal associations
- Self pollinator (wind)

[Back to Cover Crop Chart](#)



Horsehints.org

# Grain sorghum

## *(Sorghum bicolor* L. Moench)

- Warm Season, grass
- Annual
- Upright plant architecture
- Common names: Sorghum, Sudan grass,
  - *Note: These were formerly separate species that have been combined*
- Medium water use
- Fair salinity tolerance
- Seeding depth: 1 – 2 inches
- Crude protein: hay 7%, stover 5%, grain 10%
- C:N ratio: leaf 11-17, stem 10-27, root 22-30
- Benefits from arbuscular mycorrhizal associations
- Self pollinator (wind)
- Stress conditions that limit growth (e.g., drought, frost) can contribute to prussic acid accumulation in leaves



NDSU



# Corn (*Zea mays* L.)

- Warm Season, grass
- Annual
- Upright plant architecture
- High water use
- Poor salinity tolerance
- Seeding depth: 1 – 2 inches
- Crude protein: grain 9-10%, stover 5%, silage 8-11%
- C:N ratio: stalk 11-65, leaf 13-20, root 20-49
- Benefits from arbuscular mycorrhizal associations
- Self pollinator (wind)

[Back to Cover Crop Chart](#)



Midwest Cover Crops Council



USDA-ARS, NGRPL